arXiv:nucl-ex/0606005v2 2 Jun 2006

High p_T Spectra of Identified Particles Produced in Pb+Pb Collisions at 158A GeV Beam Energy

Tim Schuster and András László for the NA49 Collaboration†

C. Alt[9], T. Anticic[23], B. Baatar[8],D. Barna[4], J. Bartke[6], L. Betev[10], H. Białkowska[20], C. Blume[9], B. Boimska[20], M. Botje[1], J. Bracinik[3], R. Bramm[9], P. Bunčić[10], V. Cerny[3], P. Christakoglou[2], P. Chung[19], O. Chvala[14], J.G. Cramer[16], P. Csató[4], P. Dinkelaker[9], V. Eckardt[13], D. Flierl[9], Z. Fodor[4], P. Foka[7], V. Friese[7], J. Gál[4], M. Gaździcki[9,11], V. Genchev[18], G. Georgopoulos[2], E. Gładysz[6], K. Grebieszkow[22], S. Hegyi[4], C. Höhne[7], K. Kadija[23], A. Karev[13], D. Kikola[22], M. Kliemant[9], S. Kniege[9], V.I. Kolesnikov[8], E. Kornas[6], R. Korus[11], M. Kowalski[6], I. Kraus[7], M. Kreps[3], A. László[4], R. Lacey[19], M. van Leeuwen[1], P. Lévai[4], L. Litov[17], B. Lungwitz[9], M. Makariev[17], A.I. Malakhov[8], M. Mateev[17], G.L. Melkumov[8], C. Meurer[9], A. Mischke[1], M. Mitrovski[9], J. Molnár[4], St. Mrówczyński[11], V. Nicolic[23], G. Pálla[4], A.D. Panagiotou[2], D. Panayotov[17], A. Petridis[2], W. Peryt[22], M. Pikna[3], J. Pluta[22], D. Prindle[16], F. Pühlhofer[12], R. Renfordt[9], C. Roland[5], G. Roland[5], M. Rybczyński[11], A. Rybicki[6,10], A. Sandoval[7], N. Schmitz[13], T. Schuster[9], P. Seyboth[13], F. Siklér[4], B. Sitar[3], E. Skrzypczak[21], M. Slodkowski[22], G. Stefanek[11], R. Stock[9], C. Strabel[9], H. Ströbele[9], T. Susa[23], I. Szentpétery[4], J. Sziklai[4], M. Szuba[22], P. Szymanski[10,20], V. Trubnikov[20], D. Varga[4,10], M. Vassiliou[2], G.I. Veres[4,5], G. Vesztergombi[4], D. Vranić[7], A. Wetzler[9], Z. Włodarczyk[11] I.K. Yoo[15], J. Zimányi[4]

[1]NIKHEF, Amsterdam, Netherlands.
[2]Department of Physics, University of Athens, Athens, Greece.
[3]Comenius University, Bratislava, Slovakia.
[4]KFKI Research Institute for Particle and Nuclear Physics, Budapest, Hungary.
[5]MIT, Cambridge, USA.
[6]Institute of Nuclear Physics, Cracow, Poland.
[7]Gesellschaft für Schwerionenforschung (GSI), Darmstadt, Germany.
[8]Joint Institute for Nuclear Research, Dubna, Russia.
[9]Fachbereich Physik der Universität, Frankfurt, Germany.
[10]CERN, Geneva, Switzerland.
[11]Institute of Physics Świ etokrzyska Academy, Kielce, Poland.
[12]Fachbereich Physik der Universität, Marburg, Germany.
[13]Max-Planck-Institut für Physik, Munich, Germany.

---

† Presented at Strangeness in Quark Matter 2006, Los Angeles, California, USA

[14]Institute of Particle and Nuclear Physics, Charles University, Prague, Czech Republic.
[15]Department of Physics, Pusan National University, Pusan, Republic of Korea.
[16]Nuclear Physics Laboratory, University of Washington, Seattle, WA, USA.
[17]Atomic Physics Department, Sofia University St. Kliment Ohridski, Sofia, Bulgaria.
[18]Institute for Nuclear Research and Nuclear Energy, Sofia, Bulgaria.
[19]Chemistry Department, Stony Brook University, SUNY, Stony Brook, USA.
[20]Institute for Nuclear Studies, Warsaw, Poland.
[21]Institute for Experimental Physics, University of Warsaw, Warsaw, Poland.
[22]Faculty of Physics, Warsaw University of Technology, Warsaw, Poland.
[23]Rudjer Boskovic Institute, Zagreb, Croatia.

Abstract. Results of the NA49 collaboration on the production of hadrons with large transverse momentum in Pb+Pb collisions at 158A GeV beam energy are presented. A range up to $p_{\mathrm{T}} = 4$ GeV/c is covered. The nuclear modification factor R_{CP} is extracted for $\pi^{\pm}$, $K^{\pm}$ and p, and the baryon to meson ratios p/π^{+}, $\bar{p}/\pi^{-}$ and $\Lambda/K^{0}_{\mathrm{s}}$ are studied. All results are compared to other measurements at SPS and RHIC and to theoretical calculations.

Submitted to: *J. Phys. G: Nucl. Phys.*

1. Introduction

The features of high p_{T} hadron production at RHIC suggest the creation of a new state of matter [1]–[4]. While nuclear modification factors smaller than one at high p_{T} are interpreted as a sign for energy loss of high momentum partons in a dense strongly interacting medium, the measured high values in baryon/meson ratios can be explained by hadron production via quark coalescence. Studying these effects at the SPS can help to enhance the understanding of the underlying mechanisms. The presented results at the highest SPS beam energy of 158A GeV ($\sqrt{s_{\mathrm{NN}}} = 17.2$ GeV) exhibit similar trends as previously seen in RHIC data. They thus complement the picture obtained from numerous other observables (like e.g. the new results on Λ flow [5]) indicating that at the top SPS energy a state of matter resembling that at RHIC is created.

2. Analysis

The main tracking device of NA49 are four large volume time projection chambers (TPCs). A calorimetric measurement of the projectile spectator energy provides an independent control of the collision centrality. For a detailed description of the detector, see [6].

$\pi^{\pm}$, $K^{\pm}$, p and $\bar{p}$ are identified via their specific energy loss in the TPCs (dE/dx). The resolution of this measurement is 3–6% and the tracking efficiency is above 95%. In this analysis of charged hadrons, the collision centrality ranges (0–5)%, (12.5–23.5)% and (33.5–80)% of the total inelastic cross section were used. The presented results



Figure 1. The nuclear modification factor R_{CP} for pions, kaons and protons measured by NA49 in Pb+Pb collisions at $\sqrt{s_{\mathrm{NN}}} = 17.2$ GeV, compared to pQCD model calculations taking into account partonic energy loss (dashed line, [7]). The compared centrality ranges are indicated as fraction of the total inelastic cross section $\sigma/\sigma^{\mathrm{inel.}}_{\mathrm{tot.}}$.

refer to the rapidity interval $-0.3 < y_{\mathrm{CM}} < 0.7$. Neutral strange hadrons are identified by a reconstruction of the decay topologies $K^0_{\mathrm{s}} \rightarrow \pi^+\pi^-$ (BR $= 68.95\%$) and $\Lambda \rightarrow p\pi^-$ (63.9%). Candidates for these decays are selected by geometrical and kinematical criteria and their invariant mass is calculated to extract yields in $y - p_{\mathrm{T}}$-bins. Here, the centrality interval (0–23.5)% and the range $-0.5 < y_{\mathrm{CM}} < 0.5$ around midrapidity were used.

All results are corrected for acceptance and reconstruction inefficiency. The Λ, proton and pion yields have not yet been corrected for feeddown from the decay of heavier particles, but the bias has been estimated to be below 10%.

3. Results

The nuclear modification factor R_{CP} is defined as the yield ratio in central over peripheral collisions, scaled by the mean number of binary nucleon-nucleon collisions $\langle N_{\mathrm{coll}} \rangle$:

$$R_{\mathrm{CP}} := \frac{\langle N_{\mathrm{coll}} \rangle^{\mathrm{Per.}}}{\langle N_{\mathrm{coll}} \rangle^{\mathrm{Cen.}}} \frac{\left(\mathrm{d}^2 N / \left(\mathrm{d}p_{\mathrm{T}} \mathrm{d}y\right)\right)^{\mathrm{Cen.}}}{\left(\mathrm{d}^2 N / \left(\mathrm{d}p_{\mathrm{T}} \mathrm{d}y\right)\right)^{\mathrm{Per.}}}$$

$\langle N_{\mathrm{coll}} \rangle$ was calculated for each centrality interval using the VENUS model. The scaling may be model dependent and the corresponding systematic error in R_{CP} is assumed to be of the order of ±20%. A shaded band indicates this error in Fig. 2.

R_{CP} for charged pions, kaons and protons is shown in Fig. 1. A monotonous rise leads up to a plateau that is reached at $p_{\mathrm{T}} = 1$ GeV/c (for pions and kaons), and at $p_{\mathrm{T}} = 2$ GeV/c (for protons) respectively. A perturbative QCD calculation taking radiative energy loss of partons into account [7] is consistent with the data when considering statistical and systematic errors. Figure 2 shows a comparison to other data



Figure 2. A comparison of R_{CP} measurements in Pb+Pb collisions at the top SPS energy of $\sqrt{s_{\mathrm{NN}}} = 17.2$ GeV: π^0 from WA98 (left panel, [8]), K_{s}^0 and Λ from NA57 (middle and right panels, [9]), together with the NA49 results. The shaded bands indicate a systematical error on the NA49 results (see text).



Figure 3. The baryon/meson ratios measured in NA49 at $\sqrt{s_{\mathrm{NN}}} = 17.2$ GeV compared to results from RHIC [10],[11] at $\sqrt{s_{\mathrm{NN}}} = 200$ GeV. Please note the different scales of the plots.

available at the highest SPS energy of 158A GeV [8],[9]. When considering all systematic errors introduced by different methods of centrality determination and the calculation of $\langle N_{\mathrm{coll}} \rangle$, the results of all three experiments are consistent. R_{CP} tends towards the same values at $p_{\mathrm{T}} > 2$ GeV/c for the compared hadron species.

While at RHIC the high values for baryon/meson ratios at intermediate p_{T} can be explained by quark coalescence models (see e.g. [10]), no calculations of such models are available for SPS energies yet. The upper panels in Fig. 3 show the NA49 results for p/π^+, $\bar{p}/\pi^-$ and Λ/K_{s}^0 together with the corresponding results from PHENIX [11] and

STAR [10] at the highest RHIC energy of $\sqrt{s_{\mathrm{NN}}} = 200$ GeV. The higher net baryon density at SPS energy is manifested in quantitative differences, but the shape of the baryon/meson ratios as a function of p_{T} is the same for both energies. This can be seen in the lower panels of Fig. 3, where the double ratios are shown. New results from RHIC Cu+Cu runs at various energies also show no qualitative change with energy in the shape of the baryon/meson ratios [12].

4. Conclusions

The presented results on R_{CP} at the highest SPS energy of $\sqrt{s_{\mathrm{NN}}} = 17.2$ GeV are consistent with a pQCD model that incorporates parton energy loss and that is used to explain RHIC results. Although large systematic errors remain, the results of three different SPS experiments are in line with this picture. Baryon/meson ratios as a function of p_{T} have the same shape as at RHIC, but the theoretical interpretation in the SPS energy range is still missing.

Acknowledgments

This work was supported by the US Department of Energy Grant DE-FG03-97ER41020/A000, the Bundesministerium für Bildung und Forschung, Germany (06F137), the Virtual Institute VI-146 of Helmholtz Gemeinschaft, Germany, the Polish State Committee for Scientific Research (1 P03B 097 29, 1 PO3B 121 29, 2 P03B 04123), the Hungarian Scientific Research Foundation (T032648, T032293, T043514), the Hungarian National Science Foundation, OTKA, (F034707), the Polish-German Foundation, the Korea Research Foundation Grant (KRF-2003-070-C00015) and the Bulgarian National Science Fund (Ph-09/05).

References

[1] I. Arsene *et al.* (the BRAHMS Collaboration): Nucl. Phys. A **757** (2005) 1.
[2] B. B. Back *et al.*: Nucl. Phys. A **757** (2005) 28.
[3] J. Adams *et al.* (the STAR Collaboration): Nucl. Phys. A **757** (2005) 102.
[4] K. Adcox *et al.* (the PHENIX Collaboration): Nucl. Phys. A **757** (2005) 184.
[5] D. Kikola *et al.* (the NA49 Collaboration): these proceedings.
[6] S. V. Afanasiev *et al* (the NA49 collaboration): Nucl. Instrum. Meth. A **430** (1999), 210.
[7] X.-N. Wang: Phys. Lett. B **595** (2004), 165-170.
[8] M. M. Aggarwal *et al.* (the WA98 Collaboration): Eur. Phys. J. C **23** (2002) 225.
[9] F. Antinori *et al.* (the NA57 Collaboration), Phys. Lett. B **623** (2005) 17.
[10] J. Adams *et al.* (the STAR Collaborations): arXiv:nucl-ex/0601042.
[11] S. S. Adler *et al.* (the PHENIX Collaboration): Phys. Rev. C **69** (2004) 034909.
[12] D. Morrison *et al.* (the PHENIX Collaboration): these proceedings.